SAS Output Page 1 of 1

## Managing current weight: lose, maintain or do neither? - {Calculated Variable)\* Kentucky BRFSS 2001

Committee	T-4-1	Trying to lose weight	Maintain weight	Neither
Geographic Groups	Total Respond.#	N % C.I.(95%)	N % C.I.(95%)	N % C.I.(95%)
TOTAL	7306	2738 37.5 (35.9-39.1)	1718 24.0 (22.6-25.4)	2850 38.6 (36.9-40.2)
ADDs				
Barren River	520	183 33.0 (28.6-37.6)	147 30.1 (25.7-34.9)	190 36.9 (32.2-41.9)
Big Sandy	419	158 36.9 (31.8-42.3)	103 25.4 (20.7-30.6)	158 37.7 (32.5-43.3)
Bluegrass	518	198 35.3 (30.8-40.2)	125 25.8 (21.4-30.7)	195 38.9 (33.8-44.2)
Buffalo Trace	413	154 34.6 (26.7-43.4)	87 27.3 (15.8-42.9)	172 38.1 (30.0-47.1)
Cumberland Valley	452	173 38.4 (33.3-43.7)	98 21.8 (17.8-26.4)	181 39.8 (34.7-45.2)
FIVCO	552	214 38.2 (33.6-43.0)	124 21.8 (18.1-25.9)	214 40.0 (35.3-44.9)
Gateway	570	198 36.8 (30.7-43.3)	140 24.2 (20.0-29.1)	232 39.0 (33.8-44.5)
Green River	398	142 34.9 (29.7-40.5)	93 24.5 (19.7-30.0)	163 40.6 (35.0-46.4)
Kentucky River	466	181 36.3 (31.4-41.6)	91 21.0 (16.8-26.0)	194 42.7 (37.4-48.1)
KIPDA	547	223 40.8 (36.1-45.6)	130 21.7 (18.1-25.7)	194 37.5 (32.9-42.4)
Lake Cumberland	470	159 33.7 (28.9-38.8)	104 21.7 (17.6-26.5)	207 44.6 (39.4-49.9)
Lincoln Trail	487	186 37.1 (32.4-42.1)	108 22.7 (18.7-27.2)	193 40.2 (35.3-45.3)
Northern Kentucky	537	221 40.5 (35.9-45.3)	126 23.8 (20.0-28.2)	190 35.6 (31.1-40.5)
Pennyrile	427	173 41.4 (36.1-46.8)	105 24.9 (20.5-29.8)	149 33.8 (28.9-39.0)
Purchase	530	175 34.0 (29.5-38.8)	137 25.2 (21.3-29.4)	218 40.8 (36.2-45.7)

<sup>\*</sup>Data in this table were derived from two weight control questions regarding losing or maintaining current weight.

#Use caution in interpreting data for total respondents less than 50. N = Cell Size, NH = Non-Hispanic, %

Percentages are weighted to population characteristics.

Unless stated differently above, Missing, 'Don't know', and Refused responses are excluded.

<sup>=</sup> Percentage, C.I.(95%) = Confidence Interval (at 95 percent probability level).